

Published by: ARDC, Bajo, Wangduephodrang
PO Box No. 1263
Telephone: +975-2-481209/260 Fax: +975-2-481311
Website: www.rcbajo.gov.bt
Content by: Field Crops Sector
Contact: Mahesh Ghimiray
Email: mghimiray@gmail.com

Introduction

Mungbean (*Vigna radiata*) and Urdbean (*Vigna mungo*) are important pulse crops. Both are important short duration grain legume crops with wide adaptability, low input requirement and have the ability to improve soil fertility by fixing atmospheric nitrogen. There is often confusion between mungbean and urdbean, the two being different only at the species (*radiata* and *mungo*) level. What our farmers traditionally grow in the southern foothills is actually Urdbean, but we are used to calling it Mungbean. There are many local varieties of Urdbean but none in Mungbean. Again, there are no improved varieties of Urdbean, but there are two improved varieties of Mungbean (Bari Mung 2 and KPS-2) released from RDC Wengkharr.

Mungbean is also known as green gram and serve as a major source of dietary protein. The nutritive value of both urdbean and mungbean lies in its high and easily digestible protein, and contain approximately 25-28% protein, 1.0% oil, 3.5-4.5% fiber, 4.5-5.5% ash and 62-65% carbohydrates on dry weight basis. Urdbean is also known as black gram. The seed colour can be either black or yellow. High values of lysine make urdbean and mungbean an excellent complement to rice in terms of balanced human nutrition.

The mung bean resembles the black gram but there are differences: the corolla of *Vigna mungo* is bright yellow while that of *Vigna radiata* is pale yellow; mung bean pods are pendulous (hanging) whereas they are erect in black gram. Mung bean is slightly less hairy than black gram.

Production Techniques

Selection of Land and its Preparation

It is a common practice for our farmers to cultivate urdbean on rice bunds and terrace walls after rice crop is transplanted. It is a form of intercropping or companion cropping. This is possible in the wetland farming system. However, urdbean can be grown as a sole crop in the dryland system. It can be cultivated in different kind of soils, but the best soil for its cultivation is loam soil with good drainage. The crop should not be raised on alkaline,

saline or waterlogged soils. A well prepared seedbed is required for proper germination and establishment of the crop. For land preparation, 2-3 ploughings followed by planking or harrowing to make the seedbed free from clods and weeds will be necessary.

Selection of Cultivar/Variety

There are so far no recommended improved varieties of urdbean. The local varieties presently grown by farmers have probably been introduced from neighboring countries and selections made over the years. Generally, varieties with black seed colour and yellow seed coat are popular. For mungbean, two improved varieties from Bangladesh and Taiwan have been introduced and released in 2002 by RDC Wengkharr. These are Bari Mung 2 and KPS-2.

Seed Treatment and Seed Rate

For the prevention of soil and seed borne diseases, seeds should be treated with Rhizobium and Phosphorus Solubilising Bacteria. Seed should be treated with 2.5 g thiram or 2 g carbendazim/kg of seed. After seed treatment, the seed should be mixed with Rhizobium culture. 250g of Rhizobium culture is sufficient for the seed required for one acre. Rhizobium treatment increases the nodule formation. Generally, a seed rate of 20 - 25 kg/ha of mungbean and urdbean is used. If sown in lines, the distance between two lines should be 30 - 35 cm. Seed should be sown at a depth of 4-5 cm.

Sowing Time

Urdbean should be sown during the month of June and July. This is a common practice of our farmers who dibble seeds on bunds and terrace walls after the rice crop is transplanted. Sowing on drylands can be done at the same time, or earlier.

Fertilizers and Irrigation

Normally, urdbean is cultivated without any fertilization. However, for optimum yields, a fertilizer rate of 10 kg nitrogen, 45 kg phosphorus and 20 kg of sulphur per ha should be applied at sowing time. Use of gypsum @ 200 kg/ha would ensure availability of calcium and sulphur. It is advisable to use fertilizers on the basis of soil test and

recommendations. Irrigation is normally not required as the crop season coincides with the rainfall. However, one irrigation before flowering and another at pod-filling stage would ensure healthy seeds. Water logging in the field should be avoided at all cost.

Diseases

Both mungbean and urdbean are infested by similar fungal pathogens and viruses. Some of these diseases can cause severe losses in yield in epidemic conditions. Main diseases are described below.

Yellow Mosaic Disease

This disease is caused by yellow mosaic virus (MYMV), which is transmitted by whitefly. This viral disease is found on several alternate and collateral host which act as primary sources of inoculums. The tender leaves show yellow mosaic spots, which increase with time leading to complete yellowing. Yellowing leads to less flowering and pod development.



Early infection often leads to death of plants.

Management:

Diseased plants should be rogued out to prevent further spread of the disease. In order to prevent whitefly infestation spray with any available insecticides.

Leaf Crinkle

This disease is caused by leaf crinkle virus (ULCV) which is transmitted by aphids, whitefly and leaf hoppers. Disease symptoms include crinkling, curling, and puckering of leaves often coupled with stunting and malformation of floral organs. Enlargement in size followed by crinkle surface of laminae are the

